

# Underwater explosions and underwater noise as possible causes for harbour porpoise bycatch

**Prof. Prof. h. c. Dr. Ursula Siebert, Dr. Maria Morell, Dr. Tobias Schaffeld, Dr. Joseph Schnitzler, Dr. Jan Lakemeyer**

**Institute for Terrestrial and Aquatic Wildlife Research, Büsum, Germany**

**Dr. Julius Stürznickel, PD Dr. Ralf Oheim, Prof. Dr. Michael Ameling**

**Department of Osteology and Biomechanics, University Medical Center Hamburg-Eppendorf, Germany**

**Sven Koschinski, Meereszoologie, Germany**

# Sources of noise



Sven Koschinski



FWG, Kiel




# Impact zones of noise

Lethal injuries

Hearing impairment

Behavioural reactions


Masking






# Explosions

Sonogram of a submarine nuclear test on the Mururoa Atoll on Oct. 21<sup>st</sup> 1992: recorded at Point Sur, California



© Sound Images of the Ocean: in Research and Monitoring, Peter Wille



280 dB


Underwater sound file of a nuclear explosion at the Mururoa Atoll on 27.10.1965  
Recorded at a distance of  
**6600 km**





# Investigation of the effects of anthropogenic sounds on harbour porpoises


## Blast trauma / acoustic trauma in harbour porpoises from the Baltic Sea



- Forty-two British ground mines of the type MK 1-7 from World War II were cleared by means of blasting
- In the period from 28 August to 31 August 2019 by a NATO unit (participation of the German Navy)
- in the Exclusive Economic Zone within the MPA Fehmarnbelt, Germany

# Blast injury / acoustic trauma in harbour porpoises from the Baltic Sea

- Forensic pathological investigations including extended image processing were conducted on 24 harbour porpoises funded by the Federal Agency for Nature Conservation
- Conducted by the University of Veterinary Medicine Hannover and the University Medical Center Hamburg-Eppendorf, Germany
- 24 harbour porpoises collected through the stranding network in Schleswig-Holstein between Sept to Nov 2019



- Tympano-periotic complex of a harbour porpoise, indicating the transverse sectional planes used for histopathology.
- Decalcified tympano-periotic complex of a porpoise in cross-section.
- 3D reconstruction of a tympano-periotic complex of a harbour porpoise by HR-pQCT scan.

Photo: ITAW-TiHo

# Blast trauma / acoustic trauma in harbour porpoises from the Baltic Sea




## Summary:

### 1. Out of the 24 harbour porpoises

8 animals died of blast/acoustic trauma,

1 animal died of blast/acoustic trauma and blunt trauma

1 animal died of blast/acoustic trauma and bycatch



2. Explosions during military activities and removal of ammunition by blast is an enormous threat for harbour porpoises and other marine animals due to direct and indirect effects. Application of noise mitigation measures and deterrence of animals is critically needed without exceptions.


3. Bycatch may obscure noise impact as the cause of death.

4. Systematic investigations on the hearing system need to be allowed and conducted by specialists to understand effects of acoustic impacts.

Photo: ITAW-TiHo



# Unusual harbour porpoise mass stranding and bycatch event 2005




Wright et al. 2013. Possible Causes of a Harbour Porpoise Mass Stranding in Danish Waters in 2005. PLOSone

- 85 strandings within ~one week, 27 with signs of bycatch
- 19 post-mortem examinations, 6 reported & 3 suspicious for bycatch
- Possible contributing factors:
- Wind, currents, tides
- High season in lumpfish fisheries
- Occurrence of prey (whiting) might have attracted porpoises

# Unusual harbour porpoise mass stranding and bycatch event 2005

- NATO exercise with 85 ships, **including sonar use for mine clearance activities**
- -> Sonar exercises could have affected the attention of harbour porpoises and may have therefore led indirectly to increased bycatches
- -> hearing impairment and blast trauma was not investigated!
- It can only be speculated whether blast trauma as initial reason for death has been obscured by bycatch


Wright et al. 2013. Possible Causes of a Harbour Porpoise Mass Stranding in Danish Waters in 2005. PLOSone




Siebert & P. D. Jepson & P. Wohlsein, Eur J Wildl Res (2013) 59: 441 – 444 First indication of gas embolism in a harbour porpoise (*Phocoena phocoena*) from German waters, U.



# Pathological findings in the ear complex of bycaught animals



Haemorrhages in the tympanic cavity



Haemorrhages in the cochlea



Fracture of the TP complex with sequestration of bone fragments

Wohlsein et al., 2019. J. Comp. Pathol

Stiftung Tierärztliche Hochschule Hannover  
University of Veterinary Medicine Hannover, Foundation

Photos: ITAW-TiHo




# Underwater noise and bycatch

## Conclusions:

The contribution of underwater noise effects as causes for bycatch need to be investigated in detail in order to find successful management solutions:

- A. Cetaceans may be permanently (PTS) or temporarily (TTS) injured by underwater noise
- B. Distracted from nets by underwater noise especially young animals
- C. Showing strong behaviour reactions and being stressed by underwater noise
- D. Being unable to detect nets by masking underwater noise

Cetaceans may just be “blind” and unable to detect and react to nets and their alerting systems.



Siebert et al. 2020, Environ. Int

**Thank you for your attention**



© Bill Curtsinger